

ITP-SB-98-14
YITP-98-11
February 1998

The Determinant Representation for a Correlation Function in Scaling Lee-Yang Model

Vladimir E. Korepin¹ * and Takeshi Oota² †

¹*Institute for Theoretical Physics, State University of New York at Stony Brook,
Stony Brook, NY 11794–3840, U. S. A.*

²*Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-01, Japan*

Abstract

We consider the scaling Lee-Yang model. It corresponds to the unique perturbation of the minimal CFT model $M_{2,5}$. This is not a unitary model. We used known expression for form factors in order to obtain a closed expression for a correlation function of a trace of energy-momentum tensor. This expression is a determinant of an integral operator. Similar determinant representation were proven to be useful not only for quantum correlation functions but also in matrix models.

*e-mail: korepin@insti.physics.sunysb.edu

†e-mail: toota@yukawa.kyoto-u.ac.jp

1 Introduction

The theory of massive, relativistic, integrable models is an important part of modern quantum field theory [21]–[28]. Scattering matrices in these models factorize into a product of two-body S-matrices [21]. Form factors can be calculated on the basis of a bootstrap approach [21]–[28].

The purpose of this paper is to calculate correlation functions. As usual correlation function can be represented as an infinite series of form factors contributions. In this paper we sum up all these contributions and obtain a closed expression for the correlation function of the energy-momentum tensor. We follow an approach of the paper [5]. We introduce an auxiliary Fock space and auxiliary Bose fields (we shall call them dual fields). These fields help us to represent the form factor decomposition of a correlation function in a form similar to “free fermionic” case. This approach was developed in [29, 30, 10]. Finally a correlation function is represented as a vacuum mean value (in the auxiliary Fock space) of a determinant of an integral operator (4.3). Determinant representation were proven to be useful not only for quantum correlation functions [6] - [16] but also in matrix models [17] - [20]. It helps for asymptotical analysis of quantum correlation functions. Among other things this approach helped to calculate asymptotic of time and temperature dependent correlation function in Nonlinear Schrödinger equation [13].

The scaling Lee-Yang model can be described by the $\phi_{1,3}$ -perturbation of the non-unitary minimal model $M_{2,5}$ [1]. The theory is known to be integrable and is described by factorizable scattering theory of only one kind of particle with mass m . The two-body scattering amplitude is given by [1]

$$S(\beta) = \frac{\sinh \beta + i \sin(2\pi/3)}{\sinh \beta - i \sin(2\pi/3)}. \quad (1.1)$$

The pole at $\beta = 2\pi i/3$ corresponds to a bound state pole. The residue of the pole is negative. So three point coupling is imaginary. These facts come from non-unitarity of the theory. This S-matrix can be obtained from breather-breather S-matrix of Sine Gordon [2]. The breather-breather S-matrix of Sine Gordon was first calculated in [3].

Form factors for the trace of the energy-momentum tensor $\Theta = T_\mu^\mu/4$ is defined by the matrix element between the vacuum state $\langle \text{vac} |$ and n particle states characterized by their rapidities β_i ($i = 1, \dots, n$):

$$F_n(\beta_1, \dots, \beta_n) = \langle \text{vac} | \Theta(0) | \beta_1, \dots, \beta_n \rangle. \quad (1.2)$$

The multi-particle form factors F_n was determined in [2, 4]:

$$F_n(\beta_1, \dots, \beta_n) = H_n Q_n(x_1, \dots, x_n) \prod_{i < j} \frac{f(\beta_i - \beta_j)}{x_i + x_j}, \quad (1.3)$$

where $x_i = e^{\beta_i}$, $i = 1, \dots, n$,

$$H_n = -\frac{\pi m^2}{4\sqrt{3}} \left(\frac{i3^{1/4}}{2^{1/2}v(0)} \right)^n. \quad (1.4)$$

The function $f(\beta)$ is given by

$$f(\beta) = \frac{\cosh \beta - 1}{\cosh \beta + 1/2} v(i\pi - \beta) v(-i\pi + \beta), \quad (1.5)$$

where

$$v(\beta) = \exp \left(2 \int_0^\infty \frac{dt}{t} e^{i\beta t/\pi} \frac{\sinh(t/2) \sinh(t/3) \sinh(t/6)}{\sinh^2 t} \right). \quad (1.6)$$

The function $f(\beta)$ has a single pole in the strip $0 \leq \text{Im}\beta < \pi$ at $\beta = 2\pi i/3$ and a single zero at $\beta = 0$.

We shall use the elementary symmetric polynomials $\sigma_k(x_1, \dots, x_n)$,

$$\prod_{j=1}^n (x + x_j) = \sum_{k \in \mathbf{Z}} x^{n-k} \sigma_k(x_1, \dots, x_n),$$

It is important to remember that σ_k is equal to zero if $k < 0$ or $k > n$. The symmetric polynomials $Q_n(x_1, \dots, x_n)$ arising in the formula for form factors can be written as:

$$\begin{aligned} Q_0 &= 1, \\ Q_1 &= 1, \\ Q_2 &= \sigma_1, \\ Q_n &= \sigma_1 \sigma_{n-1} P_n, \quad n \geq 3, \end{aligned} \quad (1.7)$$

where

$$P_n = \det_{n-3} (\Sigma_{ij}). \quad (1.8)$$

Here Σ_{ij} is an $(n-3)$ by $(n-3)$ matrix with the entries

$$\Sigma_{ij} = \sigma_{3i-2j+1}, \quad 1 \leq i, j \leq n-3. \quad (1.9)$$

The index $n-3$ in the expression $\det_{n-3}$ denotes the dimension of the matrix Σ_{ij} .

After the Wick rotation to the Euclidean space, a correlation function of the operator Θ can be presented as an infinite series of form factors contributions

$$\begin{aligned} \langle \Theta(x)\Theta(0) \rangle &= \sum_{n=0}^{\infty} \int \frac{d^n \beta}{n!(2\pi)^n} \langle \text{vac} | \Theta(0) | \beta_1, \dots, \beta_n \rangle \langle \beta_n, \dots, \beta_1 | \Theta(x) | \text{vac} \rangle \\ &\times \exp \left[-mr \sum_{j=1}^n \cosh \beta_j \right], \end{aligned} \quad (1.10)$$

where $r = (x^\mu x_\mu)^{1/2}$.

In the present paper we sum up this series explicitly. Now let us present a plan of the paper. Section 2 is devoted to a transformation of the determinants to a form, which is convenient for summation. In section 3 we introduce auxiliary quantum operators—dual fields—in order to factorize an expression for a correlation function and to represent it in a form similar to “free fermionic case”. In section 4 we sum up the series (1.10) into a Fredholm determinant. In section 5 we use the Fredholm determinant representation for derivation of an asymptotic behavior of the correlation function.

2 A transformation of the form factor

A determinant of a linear integral operator $I + V$ can be written as

$$\det(I + V) = \sum_{n=0}^{\infty} \int \frac{dx_1 \cdots dx_n}{n!} \det_n \begin{pmatrix} V(x_1, x_1) & \cdots & V(x_1, x_n) \\ V(x_2, x_1) & \cdots & V(x_2, x_n) \\ \vdots & \ddots & \vdots \\ V(x_n, x_1) & \cdots & V(x_n, x_n) \end{pmatrix}. \quad (2.1)$$

Thus, in order to obtain a determinant representation for correlation functions one need to represent the form factor expansion (1.10) in the form (2.1). Determinants of integral operators, which we consider also can be called Fredholm determinants.

The form factor (1.3) is proportional to the polynomials Q_n (1.7). The polynomial Q_n for $n \geq 3$ is proportional to the determinant of the matrix Σ_{ij} (1.9).

Let M be the following n by n matrix:

$$M_{ij} = \sigma_{3i-2j-1}, \quad 1 \leq i, j \leq n. \quad (2.2)$$

Note that $M_{1j} = \delta_{1,j}$, $M_{2j} = \delta_{1,j}\sigma_3 + \delta_{2,j}\sigma_1$ and $M_{nj} = \delta_{n,j}\sigma_{n-1}$ for $j = 1, \dots, n$. Using the relation $M_{ij} = \Sigma_{i-2,j-2}$ ($i, j = 3, \dots, n-1$), it is possible to show that

$$Q_n = \det_n M, \quad n \geq 0. \quad (2.3)$$

The matrix M_{ij} consists of n^2 different functions, depending on the same set of arguments $x_1, \dots, x_n$.

The main goal of this and next section is to transform the matrix (2.2) to such a form, that entries of a new matrix would be parameterized by a single function, depending on different sets of variables, (like $V(x_i, x_j)$ in (2.1))

$$M_{ij} \rightarrow \hat{D}_{ij}, \quad \hat{D}_{ij} = \hat{D}(x_i, x_j). \quad (2.4)$$

In order to study correlation functions, we need to find a square of the polynomials:

$$Q_n^2 = \det_n (C_{ij}), \quad (2.5)$$

where

$$C_{jk} = (M^T M)_{jk} = \sum_{i=1}^n \sigma_{3i-2j-1} \sigma_{3i-2k-1} = \sum_{i=-\infty}^n \sigma_{3i-2j-1} \sigma_{3i-2k-1}. \quad (2.6)$$

Here we used the relation $\sigma_k = 0$ for $k < 0$.

Note that the elementary symmetric polynomials have the following expression:

$$\sigma_k = \frac{1}{2\pi i} \oint_{\gamma} \frac{dz}{z^{n-k+1}} \prod_{m=1}^n (z + x_m), \quad (2.7)$$

where an integration contour γ is a circle around origin in positive direction.

Substituting the above expression into (2.6) and summing up an infinite series, we have

$$C_{jk} = \oint_{\gamma_1} \frac{dz_1}{2\pi i} \oint_{\gamma_2} \frac{dz_2}{2\pi i} \frac{z_1^{2n-2j+1} z_2^{2n-2k+1}}{(z_1 z_2)^3 - 1} \prod_{m=1}^n (z_1 + x_m)(z_2 + x_m). \quad (2.8)$$

Here we have chosen the radius of the circle γ_i ($i = 1, 2$) to be greater than one in order for the series to converge.

The matrix C still depends on n^2 different functions C_{jk} . However, this matrix can be transformed to a more convenient form. Let us introduce the following matrix

$$A_{jk} = \frac{1}{(n-j)!} \frac{d^{n-j}}{d(x^2)^{n-j}} \prod_{m \neq k}^n (x^2 + x_m^2) \Big|_{x^2=0}, \quad (2.9)$$

with a determinant

$$\det A = \prod_{i < j}^n (x_i^2 - x_j^2). \quad (2.10)$$

Using this matrix, we define another matrix D , which differs from C by linear transformation

$$D = A^T C A. \quad (2.11)$$

We have an explicit expression for matrix elements of D :

$$D_{jk} = \frac{1}{(2\pi i)^2} \oint d^2 z \frac{z_1 z_2}{z_1^3 z_2^3 - 1} Y(z_1, x_j) Y(z_2, x_k), \quad (2.12)$$

where

$$Y(z, x) = \frac{J(z)}{z^2 + x^2}, \quad (2.13)$$

and

$$J(z) = \prod_{a=1}^n (z + x_a)(z^2 + x_a^2). \quad (2.14)$$

Taking the integral with respect to z_2 , we have after the symmetrization of the integrand

$$D_{jk} = \frac{1}{18\pi i} \oint_{\gamma} \frac{dz}{z} \left(\sum_{l=1}^3 \omega^l Y(\omega^{-l} z, x_j) \right) \left(\sum_{m=1}^3 \omega^m Y(\omega^{-m} z^{-1}, x_k) \right). \quad (2.15)$$

Here $\omega = e^{2\pi i/3}$ and the integration contour γ is a circle whose radius is larger than one. Explicitly, the sum in l can be written as

$$\sum_{l=1}^3 \omega^l Y(\omega^{-l} z, x) = Y(z, x) + \omega Y(\omega^{-1} z, x) + \omega^{-1} Y(\omega z, x). \quad (2.16)$$

Determinants of matrix C and D are related by

$$\det_n C = \prod_{i < j}^n (x_i^2 - x_j^2)^{-2} \det_n D. \quad (2.17)$$

Thus, we obtain a determinant representation for Q_n^2 :

$$Q_n^2 = \frac{\det_n D}{\prod_{i < j}^n (x_i^2 - x_j^2)^2}. \quad (2.18)$$

3 Dual fields

Entries of the matrix D_{jk} are parameterized now by a single function D . However, an element D_{jk} , is still not a function of two arguments only, because of the product $J(z) = \prod_{m=1}^n (z + x_m)(z^2 + x_m^2)$. This product depends on all x_m . In order to get rid of these products we introduce auxiliary Fock space and auxiliary quantum operators—dual fields.

Let us define

$$\Phi_1(x) = q_1(x) + p_2(x), \quad \Phi_2(x) = q_2(x) + p_1(x), \quad (3.1)$$

where operators $p_j(x)$ and $q_j(x)$ act in the canonical Bose Fock space in a following way

$$(0|q_j(x) = 0, \quad p_j(x)|0) = 0. \quad (3.2)$$

Non-zero commutation relations are given by

$$[p_1(x), q_1(y)] = [p_2(x), q_2(y)] = \xi(x, y) = \log((x + y)(x^2 + y^2)). \quad (3.3)$$

Due to the symmetry of the function $\xi(x, y) = \xi(y, x)$, all fields $\Phi_j(x)$ commute with each other

$$[\Phi_j(x), \Phi_k(y)] = 0, \quad j, k = 1, 2. \quad (3.4)$$

Dual fields are linear combinations of canonical Bose fields, see page 210 of [10].

Instead of $Y(z, x)$ let us define an operator valued function

$$\hat{Y}(z, x) = \frac{e^{\Phi_1(z)}}{z^2 + x^2}. \quad (3.5)$$

Instead of D_{jk} (2.15), we shall introduce an operator in the auxiliary Fock space:

$$\hat{D}(x, y) = \frac{1}{18\pi i} \oint \frac{dz}{z} \left(\sum_{l=1}^3 \omega^l \hat{Y}(\omega^{-l} z, x) \right) \left(\sum_{m=1}^3 \omega^m \hat{Y}(\omega^{-m} z^{-1}, y) \right). \quad (3.6)$$

It is easy to show that an exponent of dual field acts like a shift operator. Namely, if $g(\Phi_1(y))$ is a function of $\Phi_1(y)$ then

$$(0| \prod_{m=1}^n e^{\Phi_2(x_m)} g(\Phi_1(y)) |0) = (0| g \left(q_1(y) + \sum_{m=1}^n \xi(x_m, y) \right) |0) = g(\log J(y)).$$

Using this property of dual fields one can remove the products $J(z)$ from the matrix D_{jk} . For more detailed derivation one should look in at formula (3.6) of the paper [5].

Standard arguments of quantum field theory show that

$$\det_n D = (0| \det_n \left(\hat{D}(x_j, x_k) e^{\frac{1}{2}\Phi_2(x_j) + \frac{1}{2}\Phi_2(x_k)} \right) |0). \quad (3.7)$$

Up to now, we wrote Q_n^2 factor of $|F_n|^2$ as a determinant. An absolute value of the form factor is equal to

$$\begin{aligned} & |F_n(\beta_1, \dots, \beta_n)|^2 \\ &= |H_n|^2 Q_n^2 \prod_{i < j} \left| \frac{f(\beta_i - \beta_j)}{(x_i + x_j)} \right|^2 \\ &= |H_n|^2 \det_n D \prod_{i < j} \left| \frac{f(\beta_i - \beta_j)}{(x_i^2 - x_j^2)(x_i + x_j)} \right|^2. \end{aligned} \quad (3.8)$$

In order to factorize the double product part, we introduce another dual field

$$\tilde{\Phi}_0(x) = \tilde{q}_0(x) + \tilde{p}_0(x). \quad (3.9)$$

As usual

$$(0|\tilde{q}_0(x) = 0, \quad \tilde{p}_0(x)|0) = 0. \quad (3.10)$$

Operators $\tilde{q}_0(x)$ and $\tilde{p}_0(y)$ commute with all p_j and q_j ($j = 1, 2$). The only non-zero commutation relation is

$$[\tilde{p}_0(x), \tilde{q}_0(y)] = \eta(x, y), \quad (3.11)$$

where

$$\eta(x, y) = \eta(y, x) = 2 \log \left| \frac{f(\log \frac{x}{y})}{(x^2 - y^2)(x + y)} \right|. \quad (3.12)$$

Here we have used the fact that $|f(\beta)|$ is a symmetric function : $|f(-\beta)| = |f(\beta)|$. It is worth mentioning also that the right hand side of (3.12) has no singularity at $x = y$, because $f(\beta)$ has the first order zero at $\beta = 0$. Using some of equations for minimal form factor from [4]

$$f(\beta)f(\beta + i\pi) = \frac{\sinh \beta}{\sinh \beta - i \sin(\pi/3)}, \quad (3.13)$$

and $f(i\pi) = 4v^2(0)$, we can see that

$$f'(0) = \frac{i}{2\sqrt{3}v^2(0)}. \quad (3.14)$$

Hence

$$\eta(x, x) = -2 \log |\lambda x^3|, \quad (3.15)$$

where

$$\lambda = 8\sqrt{3}v^2(0). \quad (3.16)$$

Newly introduced dual field also commute

$$[\tilde{\Phi}_0(x), \tilde{\Phi}_0(y)] = 0 = [\tilde{\Phi}_0(x), \Phi_j(y)]. \quad (3.17)$$

Due to the Campbell–Hausdorff formula, we have

$$(0| \prod_{m=1}^n e^{\tilde{\Phi}_0(x_m)} |0) = \prod_{i,j=1}^n e^{\frac{1}{2}\eta(x_i, x_j)} = \lambda^{-n} \prod_{m=1}^n x_m^{-3} \prod_{i < j}^n \left| \frac{f(\log \frac{x_i}{x_j})}{(x_i^2 - x_j^2)(x_i + x_j)} \right|^2. \quad (3.18)$$

Combining these results, we can represent a square of an absolute value of the form factor as a determinant:

$$\begin{aligned} & |F_n(\beta_1, \dots, \beta_n)|^2 \\ &= \left(\frac{\pi m^2}{4\sqrt{3}} \right)^2 12^n \langle 0 | \det_n \left(x_j^{3/2} x_k^{3/2} \hat{D}(x_j, x_k) e^{\frac{1}{2}\Phi_0(x_j) + \frac{1}{2}\Phi_0(x_k)} \right) | 0 \rangle. \end{aligned}$$

Here

$$\Phi_0(x) = \tilde{\Phi}_0(x) + \Phi_2(x). \quad (3.19)$$

So we managed to represent a square of an absolute value of the form factor as a determinant, similar to one of the terms in the right hand side of (2.1). In the next section we shall sum up all contributions of the form factors and obtain a determinant representation for the correlation function.

4 The determinant representation

Because the scaling Lee-Yang model is a non-unitary theory, the normalization constant H_n (1.4) is pure imaginary for n odd. This leads to the following relation:

$$\langle \text{vac} | \Theta(0) | \beta_1, \dots, \beta_n \rangle \langle \beta_n, \dots, \beta_1 | \Theta(0) | \text{vac} \rangle = (-1)^n |F_n(\beta_1, \dots, \beta_n)|^2. \quad (4.1)$$

Then the correlation function of Θ can be written as

$$\begin{aligned} & \langle \Theta(x) \Theta(0) \rangle \\ &= \sum_{n=0}^{\infty} (-1)^n \int \frac{d^n \beta}{n! (2\pi)^n} |F_n(\beta_1, \dots, \beta_n)|^2 \prod_{j=1}^n e^{-\theta(x_j)} \\ &= \left(\frac{\pi m^2}{4\sqrt{3}} \right)^2 \langle 0 | \sum_{n=0}^{\infty} \int_0^{\infty} \frac{d^n x}{n!} \left(-\frac{6}{\pi} \right)^n \\ & \quad \times \det_n \left(x_j x_k \hat{D}(x_j, x_k) e^{\frac{1}{2}(\Phi_0(x_j) + \Phi_0(x_k))} e^{-\frac{1}{2}(\theta(x_j) + \theta(x_k))} \right) | 0 \rangle. \end{aligned}$$

where

$$\theta(x) = \frac{mr}{2}(x + x^{-1}). \quad (4.2)$$

Now we can use (2.1) in order to sum up. Finally we obtained a determinant representation in terms of an integral operator (Fredholm determinant)

$$\langle \Theta(x) \Theta(0) \rangle = \left(\frac{\pi m^2}{4\sqrt{3}} \right)^2 \langle 0 | \det(I - \hat{U}) | 0 \rangle. \quad (4.3)$$

The operator $\hat{U}$ act on a function $g(x)$ as

$$[\hat{U}g](x) = \int_0^\infty dy \hat{U}(x, y)g(y). \quad (4.4)$$

The kernel of the integral operator $\hat{U}(x, y)$ is equal to

$$\hat{U}(x, y) = \frac{6}{\pi} xy \hat{D}(x, y) e^{\frac{1}{2}(\Phi_0(x) + \Phi_0(y))} e^{-\frac{1}{2}(\theta(x) + \theta(y))}, \quad (4.5)$$

where $\hat{D}$ is given by (3.6)

$$\hat{D}(x, y) = \frac{1}{18\pi i} \oint \frac{dz}{z} \left(\sum_{l=1}^3 \omega^l \hat{Y}(\omega^{-l}z, x) \right) \left(\sum_{m=1}^3 \omega^m \hat{Y}(\omega^{-m}z^{-1}, y) \right), \quad (4.6)$$

where

$$\hat{Y}(z, x) = \frac{e^{\Phi_1(z)}}{z^2 + x^2}. \quad (4.7)$$

The dual fields $\Phi_0(x)$ and $\Phi_1(x)$ were defined in the section 3 (see (3.1) and (3.9)). The main property of these dual fields is, that they commute with each other, so the Fredholm determinant $\det(I - \hat{U})$ is well defined. Certainly $\det(I - \hat{U})$ is an operator in auxiliary Fock space, but it belongs to the Abelian sub algebra. On the other hand, the vacuum expectation value of these operators is non-trivial. It follows from commutation relations (3.3), (3.11), that in order to calculate the vacuum expectation value, one should use the following prescription

$$(0| \prod_{a=1}^{M_1} e^{\Phi_0(x_a)} \prod_{b=1}^{M_2} e^{\Phi_1(x_b)} |0) = \prod_{a=1}^{M_1} \prod_{b=1}^{M_1} e^{\frac{1}{2}\eta(x_a, x_b)} \prod_{a=1}^{M_1} \prod_{b=1}^{M_2} e^{\xi(x_a, x_b)}. \quad (4.8)$$

Here

$$\eta(x, y) = 2 \log \left| \frac{f(\log \frac{x}{y})}{(x^2 - y^2)(x + y)} \right|, \quad (4.9)$$

and

$$\xi(x, y) = \log((x + y)(x^2 + y^2)). \quad (4.10)$$

Similar Fredholm determinant representations were useful in the theory of correlation functions [10, 12, 15, 13].

5 Large r -asymptotic

Using the technique of [5], we calculate the long distance asymptotic of the correlation function.

The determinant can be written as

$$\det(I - \hat{U}(x, y)) = \det(I - \tilde{U}(z_1, z_2)), \quad (5.1)$$

where

$$\tilde{U}(z_1, z_2) = \int_0^\infty dx P_1(z_1, x) P_2(z_2, x), \quad (5.2)$$

and

$$P_1(z, x) = \frac{x}{3\pi^2 i z} \left(\sum_{l=1}^3 \omega^l \hat{Y}(\omega^{-l} z, x) \right) e^{\frac{1}{2}\Phi_0(x) - \frac{1}{2}\theta(x)}, \quad (5.3)$$

$$P_2(z, y) = y \left(\sum_{l=1}^3 \omega^l \hat{Y}(\omega^{-l} z^{-1}, y) \right) e^{\frac{1}{2}\Phi_0(y) - \frac{1}{2}\theta(y)}. \quad (5.4)$$

The integral operator $\tilde{U}(z_1, z_2)$ acts on a function $g(z)$ as

$$[\tilde{U}g](z_1) = \oint \tilde{U}(z_1, z_2) g(z_2) dz_2. \quad (5.5)$$

Here the integration contour is a circle around zero in positive direction.

In the limit $r \rightarrow \infty$, we evaluate the integral (5.2) by a saddle point method. The saddle point of the function $\theta(x)$ is $x = 1$. Hence, we can estimate the integral in (5.2) as

$$\tilde{U}(z_1, z_2) = P_1(z_1, 1) P_2(z_2, 1) \left(\sqrt{\frac{2\pi}{mr}} + O(r^{-3/2}) \right). \quad (5.6)$$

Thus, for the large r asymptotics the kernel $\tilde{U}(z_1, z_2)$ becomes a one-dimensional projector, and its Fredholm determinant is equal to

$$\det(I - \tilde{U}) \rightarrow 1 - \oint dz \tilde{U}(z, z). \quad (5.7)$$

Using the commutation relations of dual fields, we can evaluate $\langle 0 | \tilde{U}(z, z) | 0 \rangle$ as follows:

$$\langle 0 | \tilde{U}(z, z) | 0 \rangle = \frac{e^{-mr}}{3\pi^2 i \lambda z} \sqrt{\frac{2\pi}{mr}} Y_1(z) Y_1(z^{-1}) + \dots, \quad (5.8)$$

where

$$Y_1(z) = \sum_{l=1}^3 \omega^l (\omega^{-l} z + 1) = 3z. \quad (5.9)$$

The above result leads to the asymptotic form of the vacuum mean value of the Fredholm determinant

$$\langle 0 | \det(I - \tilde{U}) | 0 \rangle = 1 - \frac{\sqrt{3}}{2v^2(0)} \frac{e^{-mr}}{\sqrt{2\pi mr}} + \dots, \quad (5.10)$$

which agrees with the large distance contribution from zero- and one-particle states [4].

Summary

We considered scaling Lee-Yang Model and we obtained determinant representation for the correlation function of the trace of the energy-momentum tensor. We believe that determinant representations is the universal language for the description of quantum correlation functions in the integrable (exactly solvable) models of quantum field theory.

Acknowledgments

We are grateful to A. Berkovich and N. Slavnov for useful discussions. This work was supported by NSF Grant No PHY-9605226.

References

- [1] J. L. Cardy and G. Mussardo, Phys. Lett. **B 225** (1989) 275.
- [2] F. A. Smirnov, Int. J. Mod. Phys. **A4** (1989) 4213; Nucl. Phys. **B337** (1990) 156.
- [3] I. Ya. Aref'eva and V. E. Korepin, JETP Lett. **20** (1974) 312.
- [4] Al. B. Zamolodchikov, Nucl. Phys. **B348** (1991) 619.
- [5] V. E. Korepin and N. A. Slavnov, *The determinant representation for quantum correlation functions of the sinh-Gordon model*, preprint hep-th/9801046.
- [6] E. Barough, B. M. McCoy and T. T. Wu, Phys. Rev. Lett. **31** (1973) 1409;
B. M. McCoy, J. H. H. Perk and R. E. Shrock, Nucl. Phys. **B220** (1983) 35.
- [7] M. Jimbo, T. Miwa, Y. Mori and M. Sato, Physica **D1** (1990) 80.
- [8] A.G. Izergin, A.G. Pronko and N.I. Abarenkova, *Temperature correlators in the one-dimensional Hubbard model in the strong coupling limit*, preprint PDMI 5/1998, hep-th/9801167.
- [9] A. G. Izergin and A. G. Pronko, *Temperature correlators in the one-dimensional Hubbard model in the strong coupling limit*, preprint PDMI 19/1997, solv-int/9801004.
- [10] V. E. Korepin, N. M. Bogoliubov and A. G. Izergin, *Quantum Inverse Scattering Method and Correlation Functions*, Cambridge University Press (1993) p. 209.
- [11] A. R. Its, A. G. Izergin, V. E. Korepin and N. A. Slavnov, Phys. Rev. Lett. **70** (1993) 1704.
- [12] F. H. L. Eßler, H. Fhrm, A. R. Its and V. E. Korepin, J. Phys. A: Math. Gen. **29** (1996) 5619.
- [13] V. E. Korepin and N. A. Slavnov, Phys. Lett. **A236** (1997) 201.
- [14] D. Bernard and A. LeClair, Nucl. Phys. **B426** (1994) 534; Nucl. Phys. **B498** (1997) 619.
- [15] E. K. Sklyanin, *Generating function of correlators in the sl_2 Gaudin model*, preprint PDMI 10/1997, solv-int/9708007.

- [16] A. LeClair, F. Lesage, S. Sachdev and H. Saleur, Nucl. Phys. **B482** (1996) 579.
- [17] F. J. Dyson, Commun. Math. Phys. **47** (1976) 117.
- [18] C. A. Tracy and H. Widom, Commun. Math. Phys. **163** (1994) 33; Commun. Math. Phys. **179** (1996) 1, 667.
- [19] J. Harnad, C. A. Tracy and H. Widom, in *Low-Dimensional Topology and Quantum Field Theory*, ed. by H. Osborn, NATO ASI Series **B314**, Plenum Press, New York (1993) p. 231.
- [20] P. J. Forrester and A. M. Odlyzko, Phys. Rev. **E54** (1996) R4493.
- [21] A. B. Zamolodchikov and Al. B. Zamolodchikov, Ann. Phys. **120** (1979) 253;
A. B. Zamolodchikov, Advanced Studies in Pure Math. **19** (1989) 641; Int. J. Mod. Phys.**A3** (1988) 743.
- [22] B. Berg, M. Karowski and P. Weisz, Phys. Rev. **D19** (1979) 2477;
M. Karowski and P. Weisz, Nucl. Phys. **B139** (1978) 445;
M. Karowski, Phys. Rep. **49** (1979) 229.
- [23] F. A. Smirnov, *Form Factors in Completely Integrable Models of Quantum Field Theory*, World Scientific, Singapore (1992).
- [24] Al. B. Zamolodchikov, Nucl. Phys. **B348** (1991) 619.
- [25] J. L. Cardy, G. Mussardo, Nucl. Phys. **B340** (1990) 387.
- [26] S. Lukyanov, Phys. Lett. **B408** (1997) 192;
V. Brazhnikov and S. Lukyanov, *Angular quantization and form factors in massive integrable models*, preprints RU-97-58, CLNS 97/1488, hep-th/9707091.
- [27] T. Oota, Nucl. Phys. **B466** (1996) 361.
- [28] J. Balog, T. Hauer and M. R. Niedermaier, Phys. Lett. **B386** (1996) 224; Nucl. Phys. **B440** (1995) 603.
- [29] V. E. Korepin, Commun. Math. Phys. **113** (1987) 177.
- [30] N. A. Slavnov, Zap. Nauchn. Sem. POMI **245** (1997) 270.